

PROGRAM FOR IMPROVED RED COLOR

(Red or partial red varieties only)

Optional program in Nutrient Tech. Section

Three spray program

At second and third codling moth cover;

Tech Flo Copocal	1 Gal/Acre
Miracal sp	5 lbs./acre
Tech Spray Mg	1/2 pt/100 gallons

21 to 30 days prior to harvest;

Tech Spray Hi-K*	2 qts./acre
------------------	-------------

* Take a June tissue analysis. If the K level is 1.5% or higher, additional potassium is not needed. If the K level is over 2%, an aggressive calcium program should be applied to improve fruit quality and storability.

Four spray program

At second, third and fourth codling moth covers;

Tech Flo Copocal	3 Qts/Acre
Miracal sp	5 lbs./acre
Tech Spray Mg	1/2 pt/100 gallons

21 to 30 days prior to harvest;

Tech Spray Hi-K*	2 qts/acre
------------------	------------

* Take a June tissue analysis. If the K level is 1.5% or higher, additional potassium is not needed. If the K level is over 2%, an aggressive calcium program should be applied to improve fruit quality and storability.

Do not Tank mix Hi-K and Mira-Cal.

Apply at 100 to 400 gallons per acre; lower gallonage applications are believed to be most effective.

Cautions

Do not use strongly acidic materials such as Seniphos, Hi-Stop or Tri-fol in combination with, or following for up to 14 days after a Copocal. If a buffer is called for, use Tech Spray Mg @ 8 oz. per 100 gallons.

May be combined with most commonly used insecticides and fungicides.

Revised February 2008